



Botulinum Toxin Pre-Treatment Instructions

In an ideal situation it is prudent to follow some simple guidelines before treatment that can make all the difference between a fair result or great result, by reducing some possible side effects associated with the injections. We realize this is not always possible; however, minimizing these risks is always desirable.

- Patient must be in good health with no active skin infections in the areas to be treated.
- Patient should not be needle phobic.
- Avoid alcoholic beverages at least 24 hours prior to treatment. Alcohol may thin the blood which will increase the risk of bruising.
- Avoid anti-inflammatory / blood thinning medications ideally, for a period of two (2) weeks before treatment. Medications and supplements such as Aspirin, Vitamin E, Gingo Biloba, St. John's Wort, Ibuprofen, Motrin, Advil, Aleve and other NSAIDS are all blood thinning and can increase the risk of bruising/swelling after injections. Please notify our staff if you take any blood thinning prescription medication prior to scheduling an appointment.
- Schedule botulinum toxin appointment at least 2 weeks prior to a special event which may be occurring (i.e., wedding, vacation, etc.) It is not desirable to have a very special event occurring and be bruised from an injection which could have been avoided.

Ohio Institute of Osteopathic Medicine, LLC

93 North Washington Street

Tiffin, Ohio 44883

Phone: (567) 804-5777

info@oiom.org

www.oiom.org